PTO/SB/08A (04-03)

| | | - | | | | | PTO/SB/08A (04-03) | | | | |
|--|-----------------|--|------|--------|------------------------|----------------------------------|--------------------|--|--|--|--|
| _ | | MATION DI | | | Application Number | 10/670 | 0,135 | | | | |
| 🔷, | STATE | MENT BY | APPI | LICANT | Filing Date | 09/24/ | /03 | | | | |
| 1/ | Lone o | <u> </u> | | | First Named Inventor | Hong- | Ji Xu | | | | |
| MAR. | 7 3 100. | E) | | | Group Art Unit | 1661 | 1652 | | | | |
| THE STATE OF THE S | 4 | <u> </u> | | | Examiner Name | Unkno | OWA a Moore | | | | |
| Sheet | HAU | | of | 3 | Attorney Docket Number | er AND54 | 11/4-007US/58011 | | | | |
| NON-PATENT LITERATURE DOCUMENTS | | | | | | | | | | | |
| wm | C1 _, | AN, B., AND DOU, Q. P. (1996). Cleavage of retinoblastoma protein during apoptosis: an interleukin 1 converting enzyme-like protease as candidate. Cancer Res. 56, 438-442. | | | | | | | | | |
| | C2 | BABICHUK, C.K., DUGGAN, B.L., AND BLEACKLEY, R.C. (1996). In vivo regulation of murine granzyme B gene transcription in activated primary T cells. J. Biol. Chem. 271, 16485-16493. | | | | | | | | | |
| | C3 | BERTHOU, C., et al. (1995). Granzyme B and perforin lytic proteins are expressed in CD34 ⁺ peripheral blood progenitor cells mobilized by chemotherapy and granulocyte colony-stimulating factor. Blood 86, 3500-3506. | | | | | | | | | |
| | C4 , | BERTHOU, C., et al. (1997). Acquisition of granzyme B and Fas ligand proteins by human keratinocytes contributes to epidermal cell defense. J. Immunol. 159, 5293-5300. | | | | | | | | | |
| | C5 _/ | BRUNO, A.P., et al. (2000). Acute myeloblastic leukemic cells acquire cellular cytotoxicity under genotoxic stress: implication of granzyme B and perforin. Blood 96, 1914-1920. | | | | | | | | | |
| | C6 > | CHEN, P.L., et al. (1992). Stability of retinoblastoma gene expression determines the tumorigenicity of reconstituted retinoblastoma cells. Cell Growth Differ. 3, 119-125. | | | | | | | | | |
| | C7, | CHEN, W.D., et al. (1997). Apoptosis is associated with cleavage of a 5-kDa fragment from RB which mimics dephosphorylation and modulates E2F binding. Oncogene 14, 1243-1248. | | | | | | | | | |
| | C8 | CORDON-CARDO, C., AND RICHON, V.M. (1994). Expression of the retinoblastoma protein is regulated in normal human tissues. Am. J. Pathol. 144, 500-510. | | | | | | | | | |
| | C9 ′ | DYSON, N. (1998). The regulation of E2F by pRB-family proteins. Genes Dev. 12: 2245-2262. | | | | | | | | | |
| | C10, | FATTMAN, C.L., et al. (1998). p53-independent dephosphorylation and cleavage of retinoblastoma protein during tamoxifen-induced apoptosis in human breast carcinoma cells. Cancer Letters 130,103-113. | | | | | | | | | |
| | C11, | GRAUBERT, T.A., et al. (1997). Perforin/granzyme-dependent and independent mechanisms are both important for the development of graft-versus-host disease after murine bone marrow transplantation. J. Clin. Invest. 100, 904-911. | | | | | | | | | |
| C12, HAAS-KOGAN, D.A., et al. (1995). Inhibition of apoptosis by the retinoblastoma gene pro | | | | | | | | | | | |
| HEUSEL, J.W., et al. (1994). Cytotoxic lymphocytes require granzyme B for fragmentation and apoptosis in allogeneic target cells. Cell 76, 977-987. | | | | | | B for the rapid induction of DNA | | | | | |
| Examiner Signature | | W | M | Jull./ | wore | Date Considered | 28 February 2005 | | | | |

| | | | | | | | | | | | |
|--|---------------------|---|--|----------|--|----------------------|--|--|--|--|--|
| | | MATION D | | | Application Number | 10/670,135 | | | | | |
| /6 | STATE | DENT BY | APP | LICANT | Filing Date | 09/24/03 | | | | | |
| | - | ~ <u>@</u> | | | First Named Inventor | Hong-Ji Xu | | | | | |
| (M | AR 2 9 T | | | | Group Art Unit | 166+ 1652 | | | | | |
| 18 | | | | | Examiner Name | Unknown W. Moore | | | | | |
| Sheet | STRAN | | of | 3 | Attorney Docket Number | AND541/4-007US/58011 | | | | | |
| NON-PATENT LITERATURE DOCUMENTS | | | | | | | | | | | |
| www | C14 | and place | IIRST, C.E., et al. Perforin-independent expression of granzyme B and proteinase inhibitor 9 in human testis nd placenta suggests a role for granzyme B-mediated proteolysis in reproduction. Molecular Human teproduction, 7:1133-1142, 2001. | | | | | | | | |
| | C15, | HOFMANN, F., et al. (1996). The retinoblastoma gene product protects E2F-1 from degradation by the ubiquitin-proteasome pathway. Genes Dev. 10, 2949-2959. | | | | | | | | | |
| | C16 | HU, SX, et al. (2003) Expression of endogenous granzyme B in a subset of human primary breast carcinomas. Br. J. Can. 89(1):135-9. | | | | | | | | | |
| | C17 __ | HUANG, H.J., et al. (1988). Suppression of the neoplastic phenotype by replacement of the RB gene in human cancer cells. Science 242, 1563-1566. | | | | | | | | | |
| | C18 | KENNEA, NL, and MEHMET, H.J (2002) Neural stem cells, Pathol., 197(4):536-50. | | | | | | | | | |
| | C19 _, | KLEIN, J.L., et al. (1989). Genomic organization and chromosomal assignment for a serine protease gene (CSPB) expressed by human cytotoxic lymphocytes. Genomics 5, 110-117. | | | | | | | | | |
| | C20 | KONTANI, K., et al. (2001). Involvement of granzyme B and perforin in suppressing nodal metastasis of cancer cells in breast and lung cancers. Eur. J. Surg. Oncol. 27,180-186. | | | | | | | | | |
| | C21, | KREK, W., et al. (1995). Cyclin A-kinase regulation of E2F-1 DNA binding function underlies suppression of an S phase checkpoint. Cell, 83: 1149-1158. | | | | | | | | | |
| | C22 | LOBE, C.G., et al. (1986). Novel serine proteases encoded by two cytotoxic T lymphocyte-specific genes. Science 232, 858-861. | | | | | | | | | |
| | C23 | MOTYKA, B., et al. (2000) Mannose 6-phosphate/insulin-like growth factor II receptor is a death receptor for granzyme B during cytotoxic T cell-induced apoptosis. Cell 103, 491-500. | | | | | | | | | |
| | C24 _{<} | OGURA, Y., et al. (2001). Apoptosis and allograft rejection in the absence of CD8 ⁺ T cells. Transplantation 71, 1827-1834. | | | | | | | | | |
| | C25 | PINKOSKI, M. J., et al. (2000). Nuclear translocation of granzyme B in target cell apoptosis. Cell Death Differ. 7, 17-24. | | | | | | | | | |
| V | C26 | QIN, X.Q., et al. (1992). Identification of a growth suppression domain within the retinoblastoma gene product. Genes Dev. 6, 953-964. | | | | | | | | | |
| ROMANO, G., et al. (2000) Latest Developments in Gene Transfer Technology: Achievements, Perspectives, and Controversies over Therapeutic Applications Stem Cells 18: 19-39. | | | | | | | | | | | |
| Examiner Signature | | // | 60 | auso. 11 | looce Cor | e 28 Felinary 205 | | | | | |

INFORMATION DISCLOSURE **Application Number** 10/670,135 FEMENT BY APPLICANT Filing Date 09/24/03 First Named Inventor Hong-Ji Xu **Group Art Unit** 1654 1652 **Examiner Name** Unknewn More 3 AND541/4-007US/58011 Sheet of Attorney Docket Number **NON-PATENT LITERATURE DOCUMENTS C28** SCHMID, J., and WEISSMANN, C. (1987) Induction of mRNA for a serine protease and a β-thromboglobulin-like protein in mitogen-stimulated human leukocytes. J. Immunol. 139, 250-256. [JWW] SHAN, B., et al. (1996). The molecular basis of E2F-1/DP-1-induced S-phase entry and apoptosis. Cell C29 Growth Differ. 7, 689-697. SHI, L., et al. (1997). Granzyme B (GraB) autonomously crosses the cell membrane and perforin initiates C30 apoptosis and GraB nuclear localization. J. Exp. Med. 185, 855-866. SHRESTA, S., et al. (1997). Mechanisms responsible for granzyme B-independent cytotoxicity. Blood 89, C31 4085-4091. SUN, J., et al. (2001). Importance of the P4' residue in human granzyme B inhibitors and substrates revealed C32 by scanning mutagenesis of the proteinase inhibitor 9 reactive center loop. J. Biol. Chem. 276,15177-15184. TAN, X., et al. (1997). Degradation of retinoblastoma protein in tumor necrosis factor and CD95-induced C33 cell death. J. Biol. Chem. 272, 9613-9616. C34 TRAPANI, J. A., et al. (1996). Localization of granzyme B in the nucleus. A putative role in the mechanism of cytotoxic lymphocyte-mediated apoptosis. J.Biol.Chem. 271, 4127-4133. C35 WARGNIER, A., et al. (1995). Identification of human granzyme B promoter regulatory elements interacting with activated T-cell-specific proteins: implication of Ikaros and CBF binding sites in promoter activation. Proc. Natl. Acad. Sci. U. S. A. 92, 6930-6934. C36 XU, H. -J. (1995). Altered retinoblastoma (RB) protein expression in human malignancies. Adv. Anat. Pathol. 2, 213-226. XU, H.-J., et al. (1997). Reexpression of retinoblastoma protein induces tumor cell senescence and **C37** telomerase inhibition. Oncogene 15, 2589-2596. **C38** YANG, X., et al, (1998). Granzyme B mimics apical caspases. Description of a unified pathway for transactivation of executioner caspase-3 and -7. J. Biol. Chem. 273, 34278-34283. YASUKAWA, M., et al. (2000). Granule exocytosis, and not the Fas/Fas ligand system, is the main pathway C39 of cytotoxicity mediated by alloantigen-specific CD4(+) as well as CD8(+) cytotoxic T lymphocytes in humans. Blood 95, 2352-2355. C40 ZHOU, Y., et al. (1994). Further characterization of retinoblastoma gene-mediated cell growth and tumor suppression in human cancer cells. Proc. Natl. Acad. Sci. U. S. A. 91, 4165-4169. MUM Examiner William W Moore Date 8 Ft hrwy 2005 Signature

Considered